

### **Complex Numbers 3C**


1 We have  $z_1 = 1 + i$ ,  $z_2 = 2 + i$  and  $z_3 = 2 + 2i$ . The transformed triangle can be found by directly computing the transformed values  $w_1, w_2, w_3$  of  $z_1, z_2, z_3$ :




ii This transformation represents a translation by vector  $\begin{pmatrix} -3 \\ 2 \end{pmatrix}$ .



ii This transformation represents enlargement by a factor of 2 with centre (0,0).



ii This transformation represents a rotation anticlockwise through  $\frac{\pi}{2}$  and a translation by  $\begin{pmatrix} -2\\1 \end{pmatrix}$ 



1 **d** ii This transformation represents enlargement by a factor of 3 and translation by  $\begin{pmatrix} 0 \\ -2 \end{pmatrix}$ .

zTranslation  $\begin{pmatrix} -2\\ 3 \end{pmatrix}$ Translation  $\begin{pmatrix} -2\\ 3 \end{pmatrix}$  z - 2 + 3iEnlargement scale factor
4 centre (0, 0) 4(z - 2 + 3i)Hence T: w = 4(z - 2 + 3i) = 4z - 8 + 12i

2

The transformation *T* is w = 4z - 8 + 12i

Note: a = 4, b = -8 + 12i.

3 Rotation through  $\frac{\pi}{2}$  around the origin is achieved by multiplying all values in the z-plane by i. Enlargement by a scale factor of 4 is achieved by multiplying all values in the z-plane by 4.

Therefore this transformation can be written as w = 4iz.

### **SolutionBank**

# **Further Pure Mathematics Book 2**

4 z moves on a circle |z-2|=4

$$METHOD(1) \quad w = 2z - 5 + 3i$$
  

$$\Rightarrow w + 5 - 3i = 2z$$
  

$$\Rightarrow \frac{w + 5 - 3i}{2} = z$$
  

$$\Rightarrow \frac{w + 5 - 3i}{2} - 2 = z - 2$$
  

$$\Rightarrow \frac{w + 5 - 3i - 4}{2} = z - 2$$
  

$$\Rightarrow \frac{w + 1 - 3i}{2} = z - 2$$
  

$$\Rightarrow \frac{|w + 1 - 3i|}{2} = |z - 2|$$
  

$$\Rightarrow \frac{|w + 1 - 3i|}{|2|} = |z - 2|$$
  

$$\Rightarrow |w + 1 - 3i| = 2|z - 2|$$
  

$$\Rightarrow |w + 1 - 3i| = 2(4)$$
  

$$\Rightarrow |w + 1 - 3i| = 8$$
  

$$\Rightarrow |w - (-1 + 3i)| = 8$$

So the locus of w is a circle centre (-1, 3), radius 8 with equation  $(u+1)^2 + (v-3)^2 = 64$ .

METHOD (2) 
$$|z-2| = 4$$
  
z lies on a circle, centre (2, 0), radius 4  
enlargement scale factor 2, centre 0.  
2z lies on a circle, centre (4, 0), radius 8.  
translation by a translation vector  $\begin{pmatrix} -5\\ 3 \end{pmatrix}$ .  
 $w = 2z - 5 + 3i$  lies on a circle centre (-1, 3), radius 8.


So the locus of w is a circle, centre (-1, 3), radius 8 with equation  $(u+1)^2 + (v-3)^2 = 64$ .

- 5 w = z 1 + 2i
  - **a** |z-1| = 3 circle centre (1, 0) radius 3.

*METHOD*(1) |z-1| = 3 is translated by a translation vector  $\begin{pmatrix} -1 \\ 2 \end{pmatrix}$  to give a circle, centre (0, 2), radius 3, in the *w*-plane.

 $METHOD(2) \qquad w = z - 1 + 2i$  $\Rightarrow w - 2i = z - 1$  $\Rightarrow |w - 2i| = |z - 1|$  $\Rightarrow |w - 2i| = 3$ 


The locus of w is a circle, centre (0, 2), radius 3.



5 **b** 
$$\arg(z-1+i) = \frac{\pi}{4}$$
 half-line from  $(1, -1)$  at  $\frac{\pi}{4}$  with the positive real axis.


METHOD(1)  $\arg(z-1+i) = \frac{\pi}{4}$  is translated by a translation vector  $\begin{pmatrix} -1\\2 \end{pmatrix}$  to give a half-line from (0, 1) at  $\frac{\pi}{4}$  with the positive real axis. METHOD(2) w = z - 1 + 2i $\Rightarrow w + 1 - 2i = z$ So  $\arg(z-1+i) = \frac{\pi}{4}$ becomes  $\arg(w+1-2i-1+i) = \frac{\pi}{4}$  $\Rightarrow \arg(w-i) = \frac{\pi}{4}$ 

Therefore, the locus of w is a half-line from (0, 1) at  $\frac{\pi}{4}$  with the positive real axis.



c 
$$y = 2x$$
  
 $w = z - 1 + 2i$   
 $\Rightarrow z = w + 1 - 2i$   
 $\Rightarrow x + iy = u + iv + 1 - 2i$   
 $\Rightarrow x + iy = u + 1 + i(v - 2)$   
So  $y = 2x \Rightarrow v - 2 = 2(u + 1)$   
 $\Rightarrow v - 2 = 2u + 2$   
 $\Rightarrow v = 2u + 4$ 

The locus of *w* is a line with equation v = 2u + 4.



- $\mathbf{6} \quad w = \frac{1}{z}, z \neq 0$ 
  - **a** z lies on a circle, |z| = 2

$$w = \frac{1}{z}$$
  

$$\Rightarrow |w| = \left|\frac{1}{z}\right|$$
  

$$\Rightarrow |w| = \frac{|1|}{|z|}$$
  

$$\Rightarrow |w| = \frac{1}{2}$$
  
apply  $|z| = 2$ 

Therefore the locus of w is a circle, centre (0, 0), radius  $\frac{1}{2}$ , with equation  $u^2 + v^2 = \frac{1}{4}$ .

**b** z lies on the half-line,  $\arg z = \frac{\pi}{4}$   $w = \frac{1}{z} \Rightarrow wz = 1 \Rightarrow z = \frac{1}{w}$ So  $\arg z = \frac{\pi}{4}$ , becomes  $\arg\left(\frac{1}{w}\right) = \frac{\pi}{4}$   $\Rightarrow \arg(1) - \arg(w) = \frac{\pi}{4}$   $\Rightarrow -\arg w = \frac{\pi}{4}$  arg 1 = 0 $\Rightarrow \arg w = -\frac{\pi}{4}$ 

Therefore the locus of *w* is a half-line from (0, 0) at an angle of  $-\frac{\pi}{4}$  with the positive *x*-axis. The locus of *w* has equation, v = -u, u > 0, v < 0.

### **SolutionBank**

**6** c z lies on the line y = 2x + 1

$$w = \frac{1}{z} \Rightarrow wz = 1 \Rightarrow z = \frac{1}{w}.$$
  

$$\Rightarrow x + iy = \frac{1}{u + iv}$$
  

$$\Rightarrow x + iy = \frac{1}{(u + iv)} \frac{(u - iv)}{(u - iv)}$$
  

$$\Rightarrow x + iy = \frac{u - iv}{u^2 + v^2}$$
  

$$\Rightarrow x + iy = \frac{u}{u^2 + v^2} + i\left(\frac{-v}{u^2 + v^2}\right)$$
  
So  $x = \frac{u}{u^2 + v^2}$  and  $y = \frac{-v}{u^2 + v^2}$   
Hence  $y = 2x + 1$  becomes  $\frac{-v}{u^2 + v^2} = \frac{2u}{u^2 + v^2} + 1 \qquad \times (u^2 + v^2)$   

$$\Rightarrow -v = 2u + u^2 + v^2$$
  

$$\Rightarrow 0 = u^2 + 2u + v^2 + v$$
  

$$\Rightarrow (u + 1)^2 - 1 + \left(v + \frac{1}{2}\right)^2 - \frac{1}{4} = 0$$
  

$$\Rightarrow (u + 1)^2 + \left(v + \frac{1}{2}\right)^2 = \frac{5}{4}$$
  

$$\Rightarrow (u + 1)^2 + \left(v + \frac{1}{2}\right)^2 = \left(\frac{\sqrt{5}}{2}\right)^2$$

Therefore, the locus of w is a circle, centre  $\left(-1, -\frac{1}{2}\right)$ , radius  $\frac{\sqrt{5}}{2}$ , with equation

$$(u+1)^2 + \left(v + \frac{1}{2}\right)^2 = \frac{5}{4}$$

**7**  $w = z^2$ 

**a** z moves once round a circle, centre (0, 0), radius 3. The equation of the circle, |z|=3 is also r=3.

The equation of the circle can be written as  $z = 3e^{i\theta}$ 

or  $z = 3(\cos\theta + i\sin\theta)$ 

de Moivre's Theorem.

 $\Rightarrow w = z^{2} = (3(\cos \theta + i \sin \theta))^{2}$  $= 3^{2}(\cos 2\theta + i \sin 2\theta)$  $= 9(\cos 2\theta + i \sin 2\theta)$ 

So,  $w = 9(\cos 2\theta + i \sin 2\theta)$  can be written as |w| = 9

Hence, as |w| = 9 and  $\arg w = 2\theta$  then w moves twice round a circle, centre (0, 0), radius 9.

7 **b** z lies on the real-axis  $\Rightarrow y = 0$ 

So 
$$z = x + iy$$
 becomes  $z = x$  (as  $y = 0$ )  
 $\Rightarrow w = z^2 = x^2$   
 $\Rightarrow u + iv = x^2 + i(0)$   
 $\Rightarrow u = x^2$  and  $v = 0$ 

As v = 0 and  $u = x^2 \ge 0$  then w lies on the positive real-axis including the origin, 0.

c z lies on the imaginary axis  $\Rightarrow x = 0$ So z = x + iy becomes z = iy (as x = 0)  $\Rightarrow w = z^2 = (iy)^2 = -y^2$  $\Rightarrow u + iv = -y^2 + i(0)$ 

$$\Rightarrow u = -y^2 \text{ and } v = 0$$

As v = 0 and  $u = -y^2 \le 0$  then w lies on the negative real-axis including the origin, 0.

- 8 We have transformation T given by  $w = \frac{2}{i 2z}, z \neq \frac{1}{2}$ 
  - **a** i Rearrange the transformation to get an expression for *z*:

$$w = \frac{2}{i - 2z}$$
$$w(i - 2z) = 2$$
$$i - 2z = \frac{2}{w}$$
$$2z = i - \frac{2}{w}$$
$$z = \frac{i}{2} - \frac{1}{w} = \frac{iw - 2}{2w}$$

Therefore we can write 
$$|z| = \left|\frac{iw-2}{2w}\right|$$
.  
Since  $|z| = 1$  we have that:  
 $\left|\frac{iw-2}{2w}\right| = 1$   
 $|iw-2| = |2w|$   
 $|i||w+2i| = 2|w|$   
 $|w+2i| = 2|w|$ 

Write w = u + iv, substitute into the equation and square both sides: |u + iv + 2i| = 2|u + iv|  $|u + iv + 2i|^2 = 4|u + iv|^2$   $u^2 + (v + 2)^2 = 4u^2 + 4v^2$   $3u^2 + 3v^2 - 4v - 4 = 0$   $u^2 + v^2 - \frac{4}{3}v - \frac{4}{3} = 0$ Complete the square for v $u^2 + \left(v - \frac{2}{3}\right)^2 = \frac{16}{9}$ , which is the equation of a circle

ii Since 
$$u^2 + \left(v - \frac{2}{3}\right)^2 = \frac{16}{9}$$
, the circle is centred at  $\left(0, \frac{2}{3}\right)$  and has radius  $r = \frac{4}{3}$ 


8 **b** We showed that the region |z| = 1 is mapped to a circle centred at  $\left(0, \frac{2}{3}\right)$  with radius  $r = \frac{4}{3}$ 

Thus  $|z| \leq 1$  will be either a circle and its interior, or a circle and its exterior.

The easiest way to check that is to pick a point inside the circle  $|z| \leq 1$  and see where it maps to.

Pick  $z_0 = 0$  (for example). Then  $w_0 = \frac{2}{i} = -2i$ . This lies outside of the circle centred at  $\left(0, \frac{2}{3}\right)$  with

radius  $r = \frac{4}{3}$ , so we see that the region  $|z| \leq 1$  will be mapped to:



9 We want to show that the transformation T given by  $w = \frac{1}{2-z}$ ,  $z \neq 2$  transforms the circle centred at O, radius 2, |z| = 2 to a line. First, rearrange T to obtain an expression for z:

$$w(2-z) = 1$$
  

$$2-z = \frac{1}{w}$$
  

$$z = 2 - \frac{1}{w}$$
  


$$z = \frac{2w-1}{w}$$
  
As  $|z| = 2$ , we can write:  

$$2 = \frac{|2w-1|}{|w|}$$
  

$$2|w| = |2w-1|$$
  

$$|w| = |w - \frac{1}{2}|$$

This equation represents points on the perpendicular bisector of the line segment joining (0,0) and



10 We know that the transformation T is given by  $w = \frac{z - i}{z + i}, z \neq -i$ 

**a** We want to show that the circle |z - i| = 1 in z-plane is mapped to a circle in w-plane. Begin by rearranging the transformation to obtain an expression for z: w(z + i) = z - i

$$w(z+1) = z - 1$$
  

$$wz + iw = z - i$$
  

$$z(1-w) = i(w+1)$$
  

$$z = \frac{iw+i}{1-w}$$

We know that |z - i| = 1, so subtract i from both sides:

$$z - i = \frac{iw + i}{1 - w} - i$$

$$z - i = \frac{2iw}{1 - w}$$
Use  $|z - i| = 1$ :
$$1 = \frac{|2iw|}{|1 - w|}$$
Write  $w = u + iv$  and square both sides of the equation:
$$|1 - u - iv|^2 = 4|u + iv|^2$$

$$(1 - u)^2 + v^2 = 4u^2 + 4v^2$$


$$1 - 2u + u^2 = 4u^2 + 3v^2$$

$$3u^2 + 2u + 3v^2 - 1 = 0$$

$$u^2 + \frac{2}{3}u + v^2 - \frac{1}{3} = 0$$
Complete the square
$$u^2 + \frac{2}{3}u + v^2 - \frac{1}{3} = 0$$

$$\left(u + \frac{1}{3}\right)^2 + v^2 = \frac{4}{9}$$

Which represents a circle centred  $\left(-\frac{1}{3},0\right)$ , radius  $r = \frac{2}{3}$  in the *w*-plane.



b

 $T: w = \frac{3}{2-z}, z \neq 2$ 11  $\Rightarrow w(2-z) = 3$  $\Rightarrow 2w - wz = 3$  $\Rightarrow 2w = 3 + wz$  $\Rightarrow 2w - 3 = wz$  $\Rightarrow \frac{2w-3}{w} = z$  $\Rightarrow z = \frac{2w-3}{w}$  $\Rightarrow z = \frac{2(u+iv)-3}{u+iv}$  $\Rightarrow z = \frac{(2u-3) + 2iv}{u+iv}$  $\Rightarrow z = \frac{[(2u-3)+2iv]}{u+iv} \times \frac{[u-iv]}{[u-iv]}$  $\Rightarrow z = \frac{(2u-3)u - iv(2u-3) + 2iuv + 2v^2}{u^2 + v^2}$  $\Rightarrow z = \frac{2u^2 - 3u - 2uvi + 3iv + 2uvi + 2v^2}{u^2 + v^2}$  $\Rightarrow z = \frac{2u^2 - 3u + 2v^2}{u^2 + v^2} + i \left[ \frac{3v}{u^2 + v^2} \right]$ So,  $x + iy = \frac{2u^2 - 3u + 2v^2}{u^2 + v^2} + i\left[\frac{3v}{u^2 + v^2}\right]$  $\Rightarrow x = \frac{2u^2 - 3u + 2v^2}{u^2 + v^2}$ and  $y = \frac{3v}{u^2 + v^2}$ 

#### 11 (continued)

As, 
$$2y = x \Rightarrow 2\left(\frac{3v}{u^2 + v^2}\right) = \frac{2u^2 - 3u + 2v^2}{u^2 + v^2}$$
  

$$\Rightarrow \frac{6v}{u^2 + v^2} = \frac{2u^2 - 3u + 2v^2}{u^2 + v^2}$$

$$\Rightarrow 6v = 2u^2 - 3u + 2v^2$$

$$\Rightarrow 0 = 2u^2 - 3u + 2v^2 - 6v$$

$$\Rightarrow 2u^2 - 3u + 2v^2 - 6v = 0 \quad (\div 2)$$

$$\Rightarrow u^2 - \frac{3}{2}u + v^2 - 3v = 0$$

$$\Rightarrow \left(u - \frac{3}{4}\right)^2 - \frac{9}{16} + \left(v - \frac{3}{2}\right)^2 - \frac{9}{4} = 0$$

$$\Rightarrow \left(u - \frac{3}{4}\right)^2 + \left(v - \frac{3}{2}\right)^2 = \frac{9}{16} + \frac{9}{4}$$

$$\Rightarrow \left(u - \frac{3}{4}\right)^2 + \left(v - \frac{3}{2}\right)^2 = \frac{45}{16}$$

$$\Rightarrow \left(u - \frac{3}{4}\right)^2 + \left(v - \frac{3}{2}\right)^2 = \left(\frac{3\sqrt{5}}{4}\right)^2$$

The image under T of 2y = x is a circle centre  $\left(\frac{3}{4}, \frac{3}{2}\right)$ , radius  $\frac{3\sqrt{5}}{4}$ , as required.

**12** 
$$T: w = \frac{-iz+i}{z+1}, z \neq -1$$

**a** Circle with equation  $x^2 + y^2 = 1 \Longrightarrow |z| = 1$ 

$$w = \frac{-iz + i}{z + 1}$$

$$\Rightarrow w(z + 1) = -iz + i$$

$$\Rightarrow wz + w = -iz + i$$

$$\Rightarrow wz + iz = i - w$$

$$\Rightarrow z(w + i) = i - w$$

$$\Rightarrow z = \frac{i - w}{w + i}$$

$$\Rightarrow |z| = \left|\frac{i - w}{w + i}\right|$$

$$\Rightarrow |z| = \left|\frac{i - w}{w + i}\right|$$


$$\Rightarrow |z| = \frac{|i - w|}{|w + i|}$$
Applying  $|z| = 1 \Rightarrow 1 = \frac{|i - w|}{|w + i|}$ 

$$\Rightarrow |w + i| = |i - w|$$

$$\Rightarrow |w + i| = |(-1)||(w - i)|$$

$$\Rightarrow |w + i| = |w - i|$$

The image under T of  $x^2 + y^2 = 1$  is the perpendicular bisector of the line segment joining (0, -1) to (0, 1). Therefore the line l has equation v = 0. (i.e. the u-axis.)



**12 c** Circle with equation  $x^2 + y^2 = 4 \Longrightarrow |z| = 2$ 

from part **a**  

$$w = \frac{-iz + i}{z + 1}$$

$$\Rightarrow z = \frac{i - w}{w + i}$$

$$\Rightarrow |z| = \frac{|i - w|}{|w + i|}$$
Applying  $|z| = 2 \Rightarrow 2 = \frac{|i - w|}{|w + i|}$ 

$$\Rightarrow 2 |w + i | = |i - w|$$

$$\Rightarrow 2 |w + i | = |(-1)(w - i)|$$

$$\Rightarrow 2 |w + i | = |(-1)|(w - i)|$$

$$\Rightarrow 2 |w + i | = |w - i|$$

$$\Rightarrow 2 |w + i | = |w - i|$$

$$\Rightarrow 2 |u + i(v + 1)| = |u + i(v - 1)|$$

$$\Rightarrow 2^{2} |u + i(v + 1)|^{2} = |u + i(v - 1)|^{2}$$

$$\Rightarrow 4[u^{2} + (v + 1)^{2}] = u^{2} + (v - 1)^{2}$$

$$\Rightarrow 4[u^{2} + v^{2} + 2v + 1] = u^{2} + v^{2} - 2v + 1$$

$$\Rightarrow 3u^{2} + 3v^{2} + 10v + 3 = 0$$

$$\Rightarrow u^{2} + (v + \frac{5}{3})^{2} - \frac{25}{9} + 1 = 0$$

$$\Rightarrow u^{2} + (v + \frac{5}{3})^{2} = \frac{25}{9} - 1$$

$$\Rightarrow u^{2} + (v + \frac{5}{3})^{2} = \frac{16}{9}$$

$$\Rightarrow u^{2} + (v + \frac{5}{3})^{2} = \left(\frac{4}{3}\right)^{2}$$

The image under *T* of  $x^2 + y^2 = 4$  is a circle *C* with centre  $\left(0, -\frac{5}{3}\right)$ , radius  $\frac{4}{3}$ . Therefore, the equation of *C* is  $u^2 + \left(v + \frac{5}{3}\right)^2 = \frac{16}{9}$ .

**13** 
$$T: w = \frac{4z - 3i}{z - 1}, z \neq 1$$

Circle with equation |z| = 3

$$w = \frac{4z - 3i}{z - 1},$$
  

$$\Rightarrow w(z - 1) = 4z - 3i$$
  

$$\Rightarrow wz - w = 4z - 3i$$
  

$$\Rightarrow wz - 4z = w - 3i$$
  

$$\Rightarrow z(w - 4) = w - 3i$$
  

$$\Rightarrow z = \frac{w - 3i}{w - 4}$$
  

$$\Rightarrow |z| = \left|\frac{w - 3i}{w - 4}\right|$$

Applying 
$$|z|=3 \Rightarrow 3 = \frac{|w-3i|}{|w-4|}$$
  
 $\Rightarrow 3 |w-4| = |w-3i|$   
 $\Rightarrow 3 |u+iv-4| = |u+iv-3i|$   
 $\Rightarrow 3 |(u-4)+iv| = |u+i(v-3)|$   
 $\Rightarrow 3^2 |(u-4)+iv|^2 = |u+i(v-3)|^2$   
 $\Rightarrow 9[(u-4)^2+v^2] = u^2 + (v-3)^2$   
 $\Rightarrow 9[u^2 - 8u + 16 + v^2] = u^2 + v^2 - 6v + 9$   
 $\Rightarrow 9u^2 - 72u + 144 + 9v^2 = u^2 + v^2 - 6v + 9$   
 $\Rightarrow 8u^2 - 72u + 8v^2 + 6v + 144 - 9 = 0$   
 $\Rightarrow 8u^2 - 72u + 8v^2 + 6v + 135 = 0$  ( $\pm 8$ )  
 $\Rightarrow u^2 - 9u + v^2 + \frac{3}{4}v + \frac{135}{8} = 0$   
 $\Rightarrow \left(u - \frac{9}{2}\right)^2 - \frac{81}{4} + \left(v + \frac{3}{8}\right)^2 - \frac{9}{64} + \frac{135}{8} = 0$   
 $\Rightarrow \left(u - \frac{9}{2}\right)^2 + \left(v + \frac{3}{8}\right)^2 = \frac{81}{4} + \frac{9}{64} - \frac{135}{8}$   
 $\Rightarrow \left(u - \frac{9}{2}\right)^2 + \left(v + \frac{3}{8}\right)^2 = \frac{225}{64}$   
 $\Rightarrow \left(u - \frac{9}{2}\right)^2 + \left(v + \frac{3}{8}\right)^2 = \left(\frac{15}{8}\right)^2$ 

Therefore, the circle with equation |z| = 1 is mapped onto a circle *C* with centre  $\left(\frac{9}{2}, -\frac{3}{8}\right)$ , radius  $\frac{15}{8}$ .

- **14**  $T: w = \frac{1}{z+i}, z \neq -i$ 
  - **a** Real axis in the *z*-plane  $\Rightarrow y = 0$

$$w = \frac{1}{z+i}$$

$$\Rightarrow w(z+i) = 1$$

$$\Rightarrow wz + iw = 1$$

$$\Rightarrow wz = 1 - iw$$

$$\Rightarrow z = \frac{1 - i(u + iv)}{u + iv}$$

$$\Rightarrow z = \frac{1 - i(u + iv)}{u + iv}$$

$$\Rightarrow z = \frac{(1 + v)u - iu}{(u + iv)} \times \frac{(u - iv)}{(u - iv)}$$

$$\Rightarrow z = \frac{(1 + v)u - iv(1 + v) - iu^{2} - uv}{u^{2} + v^{2}}$$

$$\Rightarrow z = \frac{(1 + v)u - uv}{u^{2} + v^{2}} + \frac{i(-v(1 + v) - u^{2})}{u^{2} + v^{2}}$$

$$\Rightarrow z = \frac{u + uv - uv}{u^{2} + v^{2}} + \frac{i(-v - v^{2} - u^{2})}{u^{2} + v^{2}}$$
So  $x + iy = \frac{u}{u^{2} + v^{2}} + \frac{i(-v - v^{2} - u^{2})}{u^{2} + v^{2}}$ 

$$\Rightarrow x = \frac{u}{u^{2} + v^{2}} and y = \frac{-v - v^{2} - u^{2}}{u^{2} + v^{2}}$$
As  $y = 0, \frac{-v - v^{2} - u^{2}}{u^{2} + v^{2}} = 0$ 

$$\Rightarrow -v - v^{2} - u^{2} = 0$$

$$\Rightarrow u^{2} + v^{2} + v = 0$$

$$\Rightarrow u^{2} + \left(v + \frac{1}{2}\right)^{2} - \frac{1}{4} = 0$$

$$\Rightarrow u^{2} + \left(v + \frac{1}{2}\right)^{2} = \left(\frac{1}{2}\right)^{2}$$

#### 14 a (continued)

Therefore, the image under T of the real axis in the z-plane is a circle  $C_1$  with centre

$$\left(0,-\frac{1}{2}\right)$$
, radius  $\frac{1}{2}$ . The equation of  $C_1$  is  $u^2 + \left(v+\frac{1}{2}\right)^2 = \frac{1}{4}$ .

**b** As 
$$x = 4$$
,  $\frac{u}{u^2 + v^2} = 4$   
 $\Rightarrow u = 4(u^2 + v^2)$   
 $\Rightarrow u = 4u^2 + 4v^2$   
 $\Rightarrow 0 = 4u^2 - u + 4v^2$  (÷4)  
 $\Rightarrow 0 = u^2 - \frac{1}{4}u + v^2$   
 $\Rightarrow 0 = \left(u - \frac{1}{8}\right)^2 - \frac{1}{64} + v^2$   
 $\Rightarrow \left(u - \frac{1}{8}\right)^2 + v^2 = \frac{1}{64}$   
 $\Rightarrow \left(u - \frac{1}{8}\right)^2 + v^2 = \left(\frac{1}{8}\right)^2$ 

Therefore, the image under T of the line x = 4 is a circle  $C_2$  with centre  $\left(\frac{1}{8}, 0\right)$ , radius  $\frac{1}{8}$ . The equation of  $C_2$  is  $\left(u - \frac{1}{8}\right)^2 + v^2 = \frac{1}{64}$ .

### **SolutionBank**

# **Further Pure Mathematics Book 2**

**15**  $T: w = z + \frac{4}{z}, z \neq 0$ 

Circle with equation  $|z| = 2 \Longrightarrow x^2 + y^2 = 4$ 

$$w = z + \frac{4}{z}$$

$$\Rightarrow w = \frac{z^2 + 4}{z}$$

$$\Rightarrow w = \frac{(x + iy)^2 + 4}{x + iy}$$

$$\Rightarrow w = \frac{x^2 + 2xyi - y^2 + 4}{x + iy}$$

$$\Rightarrow w = \frac{[(x^2 - y^2 + 4) + i(2xy)]}{x + iy}$$

$$\Rightarrow w = \frac{[(x^2 - y^2 + 4) + i(2xy)]}{(x + iy)} \times \frac{(x - iy)}{(x - iy)}$$

$$\Rightarrow w = \frac{x^3 - xy^2 + 4x + 2xy^2 + i(2x^2y - x^2y + y^3 - 4y)}{x^2 + y^2}$$

$$\Rightarrow w = \left(\frac{x^3 + xy^2 + 4x}{x^2 + y^2}\right) + i\left(\frac{y^3 + x^2y - 4y}{x^2 + y^2}\right)$$

$$\Rightarrow w = \frac{x(x^2 + y^2 + 4)}{x^2 + y^2} + \frac{iy(x^2 + y^2 - 4)}{x^2 + y^2}$$
Apply  $x^2 + y^2 + 4 \Rightarrow w = \frac{x(4 + 4)}{4} + \frac{iy(4 - 4)}{4}$ 

$$\Rightarrow w = \frac{8x}{4} + \frac{iy(0)}{4}$$

$$\Rightarrow w = 2x + 0i$$

$$\Rightarrow u + iv = 2x + 0i$$

$$\Rightarrow u = 2x, v = 0$$
As  $|z| = 2 \Rightarrow -2 \leqslant x \leqslant 2$ 
So  $-4 \leqslant 2x \leqslant 4$ 
and  $-4 \leqslant u \leqslant 4$ 

Therefore the transformation T maps the points on a circle |z| = 2 in the z-plane to points in the interval [-4, 4] on the real axis in the w-plane. Hence k = 4.

**16**  $T: w = \frac{1}{z+3}, z \neq -3$ Line with equation 2x - 2y + 7 = 0 in the *z*-plane  $w = \frac{1}{\pi + 2}$  $\Rightarrow w(z+3) = 1$  $\Rightarrow wz + 3w = 1$  $\Rightarrow wz = 1 - 3w$  $\Rightarrow z = \frac{1-3w}{w}$  $\Rightarrow z = \frac{1 - 3(u + iv)}{u + iv}$  $\Rightarrow z = \frac{1 - 3u - 3iv}{u + iv}$  $\Rightarrow z = \frac{[(1-3u) - (3v)i]}{(u+iv)} \times \frac{(u-iv)}{(u-iv)}$  $\Rightarrow z = \frac{(1-3u)u - 3v^2 - iv(1-3u) - i(3uv)}{u^2 + v^2}$  $\Rightarrow z = \frac{u - 3u^2 - 3v^2}{v^2 + v^2} + \frac{i(-v + 3uv - 3uv)}{v^2 + v^2}$  $\Rightarrow z = \frac{u - 3u^2 - 3v^2}{u^2 + v^2} + \frac{i(-v)}{u^2 + v^2}$ So,  $x + iy = \frac{u - 3u^2 - 3v^2}{u^2 + v^2} + \frac{i(-v)}{u^2 + v^2}$  $\Rightarrow x = \frac{u - 3u^2 - 3v^2}{v^2 + v^2}$ and  $y = \frac{-v}{v^2 + v^2}$ As 2x - 2y + 7 = 0, then  $2\left(\frac{u-3u^2-3v^2}{u^2+v^2}\right) - 2\left(\frac{-v}{u^2+v^2}\right) + 7 = 0$  $\Rightarrow \frac{2u - 6u^2 - 6v^2}{u^2 + v^2} + \frac{2v}{u^2 + v^2} + 7 = 0 \quad (\times (u^2 + v^2))$  $\Rightarrow 2u - 6u^2 - 6v^2 + 2v + 7(u^2 + v^2) = 0$  $\Rightarrow 2u - 6u^2 - 6v^2 + 2v + 7u^2 + 7v^2 = 0$  $\Rightarrow u^2 + 2u + v^2 + 2v = 0$  $\Rightarrow (u+1)^2 - 1 + (v+1)^2 - 1 = 0$  $\Rightarrow (u+1)^2 + (v+1)^2 = 2$  $\Rightarrow (u+1)^2 + (v+1)^2 = \left(\sqrt{2}\right)^2$ 

#### 16 (continued)

Therefore the transformation T maps the line 2x-2y+7=0 in the z-plane to a circle C with centre (-1, -1), radius  $\sqrt{2}$  in the w-plane.

#### Challenge

We know that the transformation T given by w = az + b maps points  $z_1 = 0$ ,  $z_2 = 1$  and  $z_3 = 1 + i$  to  $w_1 = 2i$ ,  $w_2 = 3i$  and  $w_3 = -1 + 3i$  respectively. Substitute  $z_1, w_1$  into T to get 2i = b. Next, substitute  $z_2, w_2$  and b into T: 3i = a + 2ia = iUsing  $z_3, w_3$  we can check the result (although it is not necessary):  $az_3 + b = i(1+i) + 2i = i - 1 + 2i = -1 + 3i = w_3$ Thus T can be written as w = iz + 2i