

## Quadratics 2D

**1 a**

$$\begin{aligned}x^2 + 6x + 1 &= 0 \\x^2 + 6x &= -1 \\(x+3)^2 - 3^2 &= -1 \\(x+3)^2 &= -1 + 9 \\(x+3)^2 &= 8 \\x+3 &= \pm\sqrt{8} \\x &= -3 \pm \sqrt{8} \\x &= -3 \pm \sqrt{4 \times 2} \\x &= -3 \pm 2\sqrt{2} \\x = -3 + 2\sqrt{2} &\text{ or } x = -3 - 2\sqrt{2}\end{aligned}$$

**b**

$$\begin{aligned}x^2 + 12x + 3 &= 0 \\x^2 + 12x &= -3 \\(x+6)^2 - 6^2 &= -3 \\(x+6)^2 &= -3 + 36 \\(x+6)^2 &= 33 \\x+6 &= \pm\sqrt{33} \\x &= -6 \pm \sqrt{33} \\x = -6 + \sqrt{33} &\text{ or } x = -6 - \sqrt{33}\end{aligned}$$

**c**

$$\begin{aligned}x^2 + 4x - 2 &= 0 \\x^2 + 4x &= 2 \\(x+2)^2 - 2^2 &= 2 \\(x+2)^2 &= 2 + 4 \\(x+2)^2 &= 6 \\x+2 &= \pm\sqrt{6} \\x &= -2 \pm \sqrt{6} \\x = -2 + \sqrt{6} &\text{ or } x = -2 - \sqrt{6}\end{aligned}$$

**d**

$$\begin{aligned}x^2 - 10x &= 5 \\(x-5)^2 - 5^2 &= 5 \\(x-5)^2 &= 5 + 25 \\(x-5)^2 &= 30 \\x-5 &= \pm\sqrt{30} \\x &= 5 \pm \sqrt{30} \\x = 5 + \sqrt{30} &\text{ or } x = 5 - \sqrt{30}\end{aligned}$$

**2 a**

$$\begin{aligned}2x^2 + 6x - 3 &= 0 \\x^2 + 3x - \frac{3}{2} &= 0 \\x^2 + 3x &= \frac{3}{2} \\\left(x + \frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2 &= \frac{3}{2}\end{aligned}$$

**a**

$$\begin{aligned}\left(x + \frac{3}{2}\right)^2 &= \frac{3}{2} + \frac{9}{4} \\\left(x + \frac{3}{2}\right)^2 &= \frac{15}{4} \\x + \frac{3}{2} &= \pm\sqrt{\frac{15}{4}} \\x = -\frac{3}{2} &\pm \sqrt{\frac{15}{2}} \\x = -\frac{3}{2} + \frac{\sqrt{15}}{2} &\text{ or } x = -\frac{3}{2} - \frac{\sqrt{15}}{2}\end{aligned}$$

**b**

$$\begin{aligned}5x^2 + 8x - 2 &= 0 \\x^2 + \frac{8}{5}x - \frac{2}{5} &= 0 \\x^2 + \frac{8}{5}x &= \frac{2}{5} \\\left(x + \frac{4}{5}\right)^2 - \left(\frac{4}{5}\right)^2 &= \frac{2}{5} \\\left(x + \frac{4}{5}\right)^2 &= \frac{2}{5} + \frac{16}{25} \\\left(x + \frac{4}{5}\right)^2 &= \frac{26}{25} \\x + \frac{4}{5} &= \pm\sqrt{\frac{26}{25}} \\x = -\frac{4}{5} &\pm \frac{\sqrt{26}}{5}\end{aligned}$$

$$x = -\frac{4}{5} + \frac{\sqrt{26}}{5} \text{ or } x = -\frac{4}{5} - \frac{\sqrt{26}}{5}$$

**c**

$$\begin{aligned}4x^2 - x - 8 &= 0 \\x^2 - \frac{1}{4}x - 2 &= 0 \\x^2 - \frac{1}{4}x &= 2 \\\left(x - \frac{1}{8}\right)^2 - \left(\frac{1}{8}\right)^2 &= 2 \\\left(x - \frac{1}{8}\right)^2 &= 2 + \frac{1}{64} \\\left(x - \frac{1}{8}\right)^2 &= \frac{129}{64}\end{aligned}$$

**2 c**  $x - \frac{1}{8} = \pm \sqrt{\frac{129}{64}}$

$$x = \frac{1}{8} \pm \frac{\sqrt{129}}{8}$$

$$x = \frac{1}{8} + \frac{\sqrt{129}}{8} \text{ or } x = \frac{1}{8} - \frac{\sqrt{129}}{8}$$

**d**  $15 - 6x - 2x^2 = 0$

$$-2x^2 - 6x + 15 = 0$$

$$x^2 + 3x - \frac{15}{2} = 0$$

$$x^2 + 3x = \frac{15}{2}$$

$$\left(x + \frac{3}{2}\right)^2 - \left(\frac{3}{2}\right)^2 = \frac{15}{2}$$

$$\left(x + \frac{3}{2}\right)^2 = \frac{15}{2} + \frac{9}{4}$$

$$\left(x + \frac{3}{2}\right)^2 = \frac{39}{4}$$

$$x + \frac{3}{2} = \pm \sqrt{\frac{39}{4}}$$

$$x = -\frac{3}{2} \pm \frac{\sqrt{39}}{2}$$

$$x = -\frac{3}{2} + \frac{\sqrt{39}}{2} \text{ or } x = -\frac{3}{2} - \frac{\sqrt{39}}{2}$$

**3 a**  $x^2 - 14x + 1 = (x - 7)^2 - 7^2 + 1$   
 $= (x - 7)^2 - 49 + 1$   
 $= (x - 7)^2 - 48$

$p = -7$  and  $q = -48$

**b**  $x^2 - 14x + 1 = 0$

$$(x - 7)^2 - 48 = 0$$

$$(x - 7)^2 = 48$$

$$x - 7 = \pm \sqrt{48}$$

$$x = 7 \pm \sqrt{16 \times 3}$$

$$x = 7 \pm 4\sqrt{3}$$

$r = 7$  and  $s = 4$

**4**  $x^2 + 2bx + c = 0$

$$(x + b)^2 - b^2 + c = 0$$

$$(x + b)^2 = b^2 - c$$

$$x + b = \pm \sqrt{b^2 - c}$$

$$x = -b \pm \sqrt{b^2 - c}$$

## Challenge

**a**  $ax^2 + 2bx + c = 0$

$$x^2 + \frac{2b}{a}x + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{a}\right)^2 - \left(\frac{b}{a}\right)^2 + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{a}\right)^2 - \frac{b^2}{a^2} + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{a}\right)^2 = \frac{b^2}{a^2} - \frac{c}{a}$$

$$\left(x + \frac{b}{a}\right)^2 = \frac{b^2 - ac}{a^2}$$

$$x + \frac{b}{a} = \pm \sqrt{\frac{b^2 - ac}{a^2}}$$

$$x = -\frac{b}{a} \pm \sqrt{\frac{b^2 - ac}{a^2}}$$

**b**  $ax^2 + bx + c = 0$

$$x^2 + \frac{b}{a}x + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2 + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{2a}\right)^2 - \frac{b^2}{4a^2} + \frac{c}{a} = 0$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2}{4a^2} - \frac{c}{a}$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2}$$

$$x + \frac{b}{2a} = \pm \sqrt{\frac{b^2 - 4ac}{4a^2}}$$

$$x = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$