


Radians Mixed exercise
1


- a** In the right-angled triangle ABC :

$$\cos \angle BAC = \frac{BA}{AC} = \frac{5}{10} = \frac{1}{2}$$

$$\text{so } \angle BAC = \frac{\pi}{3}$$

- b** Area of triangle ABC


$$\begin{aligned} &= \frac{1}{2} \times AB \times AC \times \sin \angle BAC \\ &= \frac{1}{2} \times 5 \times 10 \times \sin \frac{\pi}{3} = 21.650\ldots \text{cm}^2 \end{aligned}$$

Area of sector DAB

$$= \frac{1}{2} \times 5^2 \times \frac{\pi}{3} = 13.089\ldots \text{cm}^2$$

Area of shaded region

$$\begin{aligned} &= \text{area of } \triangle ABC - \text{area of sector } DAB \\ &= 21.650\ldots - 13.089\ldots = 8.56 \text{ cm}^2 \text{ (3 s.f.)} \end{aligned}$$

2


- a** Using Pythagoras' theorem to find OM :

$$OM^2 = 17^2 - 15^2 = 64 \Rightarrow OM = 8 \text{ cm}$$

$$\begin{aligned} \text{Area of } \triangle OCD &= \frac{1}{2} \times CD \times OM \\ &= \frac{1}{2} \times 30 \times 8 = 120 \text{ cm}^2 \end{aligned}$$

- b** Area of shaded region R

$$\begin{aligned} &= \text{area of semicircle } CDA_1 \\ &\quad - \text{area of segment } CDA_2 \end{aligned}$$

$$\text{Area of semicircle } CDA_1$$

$$= \frac{1}{2} \times \pi \times 15^2 = 353.429\ldots \text{cm}^2$$

$$\text{Area of segment } CDA_2$$

$$= \text{area of sector } OCD$$

$$- \text{area of triangle } OCD$$

$$= \frac{1}{2} \times 17^2 \times \angle COD - 120$$

In right-angled triangle COD :

$$\sin \angle COM = \frac{CM}{OC} = \frac{15}{17}$$

$$\text{so } \angle COM = 1.0808\ldots$$

$$\text{hence } \angle COD = 2.1616\ldots$$

$$\text{So area of segment } CDA_2$$

$$= \frac{1}{2} \times 17^2 \times 2.1616\ldots - 120$$


$$= 192.362\ldots \text{cm}^2$$

$$\text{So area of shaded region } R$$

$$= 353.429\ldots - 192.362\ldots$$

$$= 161.07 \text{ cm}^2 \text{ (2 d.p.)}$$

3




- a Reflex angle $AOB = (2\pi - \theta)$ rad
Area of shaded sector

$$\begin{aligned} &= \frac{1}{2} \times 6^2 \times (2\pi - \theta) \\ &= (36\pi - 18\theta) \text{ cm}^2 \\ \text{So } 80 &= 36\pi - 18\theta \\ \Rightarrow 18\theta &= 36\pi - 80 \\ \Rightarrow \theta &= \frac{36\pi - 80}{18} = 1.839 \text{ (3 d.p.)} \end{aligned}$$

- b Length of minor arc AB
 $= 6\theta = 6 \times 1.8387\dots = 11.03 \text{ cm (2 d.p.)}$

4



- a Using $l = r\theta$:

$$p = r\theta \Rightarrow \theta = \frac{p}{r}$$

- b Area of sector

$$= \frac{1}{2}r^2\theta = \frac{1}{2}r^2 \times \frac{p}{r} = \frac{1}{2}pr \text{ cm}^2$$

- c $4.65 \leq r < 4.75, 5.25 \leq p < 5.35$
Least possible value for area of sector

$$= \frac{1}{2} \times 5.25 \times 4.65 = 12.207 \text{ cm}^2 \text{ (3 d.p.)}$$

(Note: Least possible value is 12.20625, so 12.207 should be given, not 12.206)

- d Maximum possible value of θ

$$= \frac{\max p}{\min r} = \frac{5.35}{4.65} = 1.1505\dots$$


So give 1.150 (3 d.p.)

Minimum possible value of θ

$$= \frac{\min p}{\max r} = \frac{5.25}{4.75} = 1.1052\dots$$

So give 1.106 (3 d.p.)

5



- a Using $l = r\theta$:

$$6.4 = 5\theta \Rightarrow \theta = \frac{6.4}{5} = 1.28 \text{ rad}$$

- b Using area of sector $= \frac{1}{2}r^2\theta$:

$$R_1 = \frac{1}{2} \times 5^2 \times 1.28 = 16$$


- c $R_2 = \text{area of circle} - R_1$

$$= \pi \times 5^2 - 16 = 62.5398\dots$$


$$\text{So } \frac{R_1}{R_2} = \frac{16}{62.5398\dots} = \frac{1}{3.908\dots} = \frac{1}{p}$$

$$\Rightarrow p = 3.91 \text{ (3 s.f.)}$$

6




Shape X



Shape Y

7



- a** Area of shape X

$$\begin{aligned} &= \text{area of rectangle} + \text{area of semicircle} \\ &= (2d^2 + \frac{1}{2}\pi d^2) \text{ cm}^2 \end{aligned}$$

$$\text{Area of shape } Y = \frac{1}{2}(2d)^2\theta = 2d^2\theta \text{ cm}^2$$

Since $X = Y$:

$$2d^2 + \frac{1}{2}\pi d^2 = 2d^2\theta$$

Divide by $2d^2$:

$$1 + \frac{\pi}{4} = \theta$$

- b** Perimeter of shape X

$$\begin{aligned} &= (d + 2d + d + \pi d) \text{ cm} \text{ with } d = 3 \\ &= (3\pi + 12) \text{ cm} \end{aligned}$$

- c** Perimeter of shape Y

$$= (2d + 2d + 2d\theta) \text{ cm}$$

$$\text{with } d = 3 \text{ and } \theta = 1 + \frac{\pi}{4}$$

$$= 12 + 6\left(1 + \frac{\pi}{4}\right)$$

$$= \left(18 + \frac{3\pi}{2}\right) \text{ cm}$$

- d** Difference

$$= \left(18 + \frac{3\pi}{2}\right) - (3\pi + 12)$$

$$= 6 - \frac{3\pi}{2}$$

$$= 1.287\ldots \text{ cm}$$

$$= 12.9 \text{ mm (3 s.f.)}$$

- a** Area of segment R_1

$$\begin{aligned} &= \text{area of sector } OPQ \\ &\quad - \text{area of triangle } OPQ \end{aligned}$$

$$\begin{aligned} \Rightarrow A_1 &= \frac{1}{2} \times 6^2 \times \theta - \frac{1}{2} \times 6^2 \times \sin \theta \\ \Rightarrow A_1 &= 18(\theta - \sin \theta) \end{aligned}$$

- b** $A_2 = \text{area of circle} - A_1$

$$\begin{aligned} &= \pi \times 6^2 - 18(\theta - \sin \theta) \\ &= 36\pi - 18(\theta - \sin \theta) \end{aligned}$$

Since $A_2 = 3A_1$:

$$36\pi - 18(\theta - \sin \theta) = 3 \times 18(\theta - \sin \theta)$$


$$36\pi - 18(\theta - \sin \theta) = 54(\theta - \sin \theta)$$

$$36\pi = 72(\theta - \sin \theta)$$

$$\pi = \theta - \sin \theta$$

$$\sin \theta = \theta - \frac{\pi}{2}$$

8



- a Using the cosine rule in $\triangle ABC$:

$$\begin{aligned}\cos A &= \frac{b^2 + c^2 - a^2}{2bc} \\ \Rightarrow \cos \angle BAC &= \frac{5^2 + 9^2 - 10^2}{2 \times 5 \times 9} = 0.06 \\ \Rightarrow \angle BAC &= 1.50408\dots \\ &= 1.504 \text{ rad (3 d.p.)}\end{aligned}$$


- b i Using the sector area formula:

$$\begin{aligned}\text{area of sector} &= \frac{1}{2}r^2\theta \\ \Rightarrow \text{area of sector } APQ &= \frac{1}{2} \times 3^2 \times 1.504 = 6.77 \text{ cm}^2 \text{ (3 s.f.)}\end{aligned}$$

ii Area of shaded region $BPQC$
 $= \text{area of } \triangle ABC - \text{area of sector } APQ$
 $= \frac{1}{2} \times 5 \times 9 \times \sin 1.504 - \frac{1}{2} \times 3^2 \times 1.504$
 $= 15.681\dots$
 $= 15.7 \text{ cm}^2 \text{ (3 s.f.)}$

iii Perimeter of shaded region $BPQC$
 $= QC + CB + BP + \text{arc length } PQ$
 $= 2 + 10 + 6 + (3 \times 1.504)$
 $= 22.51\dots$
 $= 22.5 \text{ cm (3 s.f.)}$

9



a Area of sector $= \frac{1}{2}r^2\theta = \frac{1}{2}r^2 \times 1.5 \text{ cm}^2$


$$\begin{aligned}\text{So } \frac{3}{4}r^2 &= 15 \\ \Rightarrow r^2 &= \frac{60}{3} = 20 \\ \Rightarrow r &= \sqrt{20} = \sqrt{4 \times 5} = \sqrt{4} \times \sqrt{5} = 2\sqrt{5}\end{aligned}$$

b Arc length $AB = r(1.5) = 3\sqrt{5} \text{ cm}$

$$\begin{aligned}\text{Perimeter of sector} &= AO + OB + \text{arc length } AB \\ &= 2\sqrt{5} + 2\sqrt{5} + 3\sqrt{5} \\ &= 7\sqrt{5} \\ &= 15.7 \text{ cm (3 s.f.)}\end{aligned}$$

c Area of segment R
 $= \text{area of sector} - \text{area of } \triangle AOB$
 $= 15 - \frac{1}{2}r^2 \sin 1.5$
 $= 15 - 10 \sin 1.5$
 $= 5.025 \text{ cm}^2 \text{ (3 d.p.)}$

10



- a Using the right-angled $\triangle ABD$, with

$$\angle ABD = \frac{\pi}{3} :$$

$$\sin \frac{\pi}{3} = \frac{3}{AB}$$

$$\Rightarrow AB = \frac{3}{\sin \frac{\pi}{3}} = \frac{3}{\frac{\sqrt{3}}{2}}$$

$$= 3 \times \frac{2}{\sqrt{3}} = 2\sqrt{3} \text{ cm}$$

- b Area of badge = area of sector

$$= \frac{1}{2} \times (2\sqrt{3})^2 \theta \text{ where } \theta = \frac{\pi}{3}$$

$$= \frac{1}{2} \times 4 \times 3 \times \frac{\pi}{3}$$

$$= 2\pi \text{ cm}^2$$

- c Perimeter of badge


$$= AB + AC + \text{arc length } BC$$

$$= 2\sqrt{3} + 2\sqrt{3} + 2\sqrt{3} \times \frac{\pi}{3}$$

$$= 2\sqrt{3} \left(2 + \frac{\pi}{3} \right)$$

$$= \frac{2\sqrt{3}}{3} (6 + \pi) \text{ cm}$$

11



- a Using the right-angled $\triangle ACD$:

$$\sin \angle ACD = \frac{35}{44}$$

$$\text{So } \angle ACD = \sin^{-1} \frac{35}{44}$$

$$\text{and } \angle ACB = 2 \sin^{-1} \frac{35}{44}$$

$$\Rightarrow \angle ACB = 1.8395\dots$$

$$= 1.84 \text{ rad (2 d.p.)}$$

- b i Length of railway track

$$= \text{length of arc } AB$$

$$= 44 \times 1.8395\dots$$

$$= 80.9 \text{ m (3 s.f.)}$$

- ii Shortest distance from C to AB is DC. Using Pythagoras' theorem:

$$DC^2 = 44^2 - 35^2$$

$$DC = \sqrt{44^2 - 35^2} = 26.7 \text{ m (3 s.f.)}$$

- iii Area of region


$$= \text{area of segment}$$

$$= \text{area of sector } ABC - \text{area of } \triangle ABC$$

$$= \frac{1}{2} \times 44^2 \times 1.8395\dots - \frac{1}{2} \times 70 \times DC$$

$$= 847 \text{ m}^2 \text{ (3 s.f.)}$$

12



- a In right-angled $\triangle OAX$ (see diagram):

$$\frac{x}{6} = \sin \theta$$

$$\Rightarrow x = 6 \sin \theta$$

$$\text{So } AB = 2x = 12 \sin \theta \quad (AB = DC)$$

Perimeter of the cross-section

$$= \text{arc length } AB + AD + DC + BC$$

$$= 6 \times 2\theta + 4 + 12 \sin \theta + 4 \text{ cm}$$

$$= (8 + 12\theta + 12 \sin \theta) \text{ cm}$$

$$\text{So } 2(7 + \pi) = 8 + 12\theta + 12 \sin \theta$$

$$\Rightarrow 14 + 2\pi = 8 + 12\theta + 12 \sin \theta$$

$$\Rightarrow 12\theta + 12 \sin \theta - 6 = 2\pi$$


Divide by 6:

$$2\theta + 2 \sin \theta - 1 = \frac{\pi}{3}$$

- b When $\theta = \frac{\pi}{6}$,

$$\begin{aligned} 2\theta + 2 \sin \theta - 1 &= \frac{\pi}{3} + \left(2 \times \frac{1}{2}\right) - 1 \\ &= \frac{\pi}{3} \end{aligned}$$

c




Area of the cross-section
= area of rectangle $ABCD$
– area of shaded segment

$$\begin{aligned} \text{Area of rectangle} &= 4 \times 12 \times \sin \frac{\pi}{6} \\ &= 24 \text{ cm}^2 \end{aligned}$$

$$\begin{aligned} \text{Area of shaded segment} &= \text{area of sector} - \text{area of triangle} \\ &= \frac{1}{2} \times 6^2 \times \frac{\pi}{3} - \frac{1}{2} \times 6^2 \times \sin \frac{\pi}{3} \\ &= 3.261\dots \text{cm}^2 \\ \text{So area of cross-section} &= 20.7 \text{ cm}^2 \quad (3 \text{ s.f.}) \end{aligned}$$

13



- a $O_1A = O_2A = 12$, as they are radii of their respective circles.
 $O_1O_2 = 12$, as O_2 is on the circumference of C_1 and hence is a radius (and vice versa).
 Therefore $\triangle AO_1O_2$ is equilateral.

$$\text{So } \angle AO_1O_2 = \frac{\pi}{3}$$

$$\text{and } \angle AO_1B = 2 \times \angle AO_1O_2 = \frac{2\pi}{3}$$

- 13 b** Consider arc AO_2B of circle C_1 .

Using arc length = $r\theta$:

$$\text{arc length } AO_2B = 12 \times \frac{2\pi}{3} = 8\pi \text{ cm}$$

Perimeter of R

$$\begin{aligned} &= \text{arc length } AO_2B + \text{arc length } AO_1B \\ &= 2 \times 8\pi = 16\pi \text{ cm} \end{aligned}$$

- c** Consider the segment AO_2B in circle C_1 .

Area of segment AO_2B

$$\begin{aligned} &= \text{area of sector } O_1AB - \text{area of } \triangle O_1AB \\ &= \frac{1}{2} \times 12^2 \times \frac{2\pi}{3} - \frac{1}{2} \times 12^2 \times \sin \frac{2\pi}{3} \end{aligned}$$

$$= 88.442\dots \text{ cm}^2$$

Area of region R

$$\begin{aligned} &= \text{area of segment } AO_2B \\ &\quad + \text{area of segment } AO_1B \\ &= 2 \times 88.442\dots \\ &= 177 \text{ cm}^2 \text{ (3 s.f.)} \end{aligned}$$

- 14 a** The student has used an angle measured in degrees – it needs to be measured in radians to use that formula.

b $50^\circ = \frac{50}{180} \times \pi \text{ rad}$

$$\begin{aligned} \frac{1}{2}r^2\theta &= \frac{1}{2} \times 3^2 \times \frac{5}{18}\pi \\ &= \frac{5}{4}\pi \text{ cm}^2 \end{aligned}$$

15 a $\frac{\cos \theta - 1}{\theta \tan 2\theta} \approx \frac{\left(1 - \frac{\theta^2}{2}\right) - 1}{\theta \times 2\theta}$

$$\begin{aligned} &= \frac{-\frac{\theta^2}{2}}{\theta \times 2\theta} \\ &= \frac{-\theta^2}{4\theta^2} \\ &= -\frac{1}{4} \end{aligned}$$

b

$$\begin{aligned} \frac{2(1 - \cos \theta) - 1}{\tan \theta - 1} &\approx \frac{2\left(1 - \left(1 - \frac{\theta^2}{2}\right)\right) - 1}{\theta - 1} \\ &= \frac{2 \times \frac{\theta^2}{2} - 1}{\theta - 1} \\ &= \frac{\theta^2 - 1}{\theta - 1} \\ &= \frac{(\theta - 1)(\theta + 1)}{\theta - 1} \\ &= \theta + 1 \end{aligned}$$

16 a $\frac{7 + 2 \cos 2\theta}{\tan 2\theta + 3} \approx \frac{7 + 2\left(1 - \frac{(2\theta)^2}{2}\right)}{2\theta + 3}$

$$\begin{aligned} &= \frac{7 + 2\left(1 - \frac{4\theta^2}{2}\right)}{2\theta + 3} \\ &= \frac{9 - 4\theta^2}{2\theta + 3} \\ &= \frac{(3 + 2\theta)(3 - 2\theta)}{2\theta + 3} \\ &= 3 - 2\theta \end{aligned}$$

b 3

- 17 a** When θ is small:

$$\text{LHS} = 32 \cos 5\theta + 203 \tan 10\theta$$

$$\begin{aligned} &\approx 32\left(1 - \frac{(5\theta)^2}{2}\right) + 203(10\theta) \\ &= 32 - 16(25\theta^2) + 2010\theta \end{aligned}$$

$$\text{So } 32 - 400\theta^2 - 2030\theta = 182$$

$$400\theta^2 + 2030\theta + 150 = 0$$

$$40\theta^2 + 203\theta + 15 = 0$$

- b** $40\theta^2 + 203\theta + 15 = 0$

$$(40\theta + 3)(\theta + 5) = 0$$

$$\theta = \frac{3}{40}, 5$$

17 c $\theta = 5$ is not a valid solution, as 5 is not ‘small’. $\frac{3}{40}$ is ‘small’, so this solution is valid.

$$\begin{aligned} \mathbf{18} \quad & \cos^4 \theta - \sin^4 \theta \\ &= (\cos^2 \theta - \sin^2 \theta)(\cos^2 \theta + \sin^2 \theta) \\ &= \cos^2 \theta - \sin^2 \theta \\ &= 1 - \sin^2 \theta - \sin^2 \theta \\ &= 1 - 2 \sin^2 \theta \\ &\approx 1 - 2\theta^2 \end{aligned}$$

19 a $3 \sin \theta = 2, 0 \leq \theta \leq \pi$

$$\begin{aligned} \sin \theta &= \frac{2}{3} \\ \theta &= 0.730, 2.41 \end{aligned}$$

b $\sin \theta = -\cos \theta, -\pi \leq \theta \leq \pi$

$$\tan \theta = -1$$

$$\theta = -\frac{\pi}{4}, \frac{3\pi}{4}$$

c $\tan \theta + \frac{1}{\tan \theta} = 2, 0 \leq \theta \leq 2\pi$

$$\tan^2 \theta + 1 = 2 \tan \theta$$

$$\tan^2 \theta - 2 \tan \theta + 1 = 0$$

$$(\tan \theta - 1)^2 = 0$$

$$\tan \theta = 1$$

$$\theta = \frac{\pi}{4}, \frac{5\pi}{4}$$

d $2 \sin^2 \theta - \sin \theta - 1 = \sin^2 \theta, -\pi \leq \theta \leq \pi$

$$\sin^2 \theta - \sin \theta - 1 = 0$$


$$\sin \theta = \frac{1 \pm \sqrt{5}}{2}$$

$$\sin \theta = 1.618 \text{ (no solutions)}$$

$$\text{or } \sin \theta = -0.618$$

$$\Rightarrow \theta = -0.666, -2.48$$

20 a



b The curves intersect twice in the given range, so the equation has two solutions.

c $5 \sin x = 3 \cos x$

$$\tan x = \frac{3}{5}$$

$$x = 0.540, 3.68$$

21 a $4 \sin \theta - \cos\left(\frac{\pi}{2} - \theta\right) = 4 \sin \theta - \sin \theta$
 $= 3 \sin \theta$

b $4 \sin \theta - \cos\left(\frac{\pi}{2} - \theta\right) = 1, 0 \leq \theta \leq 2\pi$
 $3 \sin \theta = 1$

$$\sin \theta = \frac{1}{3}$$

$$\theta = 0.340, 2.80$$

22 $\frac{\sin 2x + 0.5}{1 - \sin 2x} = 2, 0 < x < \frac{3\pi}{2}$
 $\sin 2x + 0.5 = 2 - 2 \sin 2x$

$$3 \sin 2x = 1.5$$

$$\sin 2x = 0.5$$

$$\text{Let } X = 2x$$

$$\sin X = 0.5, 0 < X < 3\pi$$

$$X = \frac{\pi}{6}, \frac{5\pi}{6}, \frac{13\pi}{6}, \frac{17\pi}{6}$$

$$x = \frac{\pi}{12}, \frac{5\pi}{12}, \frac{13\pi}{12}, \frac{17\pi}{12}$$

23 a Cosine can be negative, so do not reject

$$-\frac{1}{\sqrt{2}}$$

b Rearranged incorrectly, so incorrectly square rooted.

c $2 \cos^2 x = 1$

$$\cos^2 x = \frac{1}{2}$$

$$\cos x = \pm \frac{1}{\sqrt{2}}$$

$$x = -\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{\pi}{4}, \frac{3\pi}{4}$$

24 a Not all solutions have been calculated. There will be four solutions in the given interval.

b $2 \tan 2x = 5, 0 \leq x \leq 2\pi$

$$\text{Let } X = 2x$$

$$2 \tan X = 5, 0 \leq X \leq 4\pi$$

$$\tan X = 2.5$$

$$X = 1.19, 4.33, 7.47, 10.6$$

$$x = 0.595, 2.17, 3.74, 5.31$$

25 a $5 \sin x = 1 + \cos^2 x$

$$5 \sin x = 1 + 2(1 - \sin^2 x)$$

$$2 \sin^2 x + 5 \sin x - 3 = 0$$

b $2 \sin^2 x + 5 \sin x - 3 = 0, 0 \leq x \leq 2\pi$

$$(2 \sin x - 1)(\sin x + 3) = 0$$

$$\sin x = 0.5 \Rightarrow x = \frac{\pi}{6}, \frac{5\pi}{6}$$

or $\sin x = -3$ (no solution)

26 a $4 \sin^2 x + 9 \cos x - 6 = 0$

$$4(1 - \cos^2 x) + 9 \cos x - 6 = 0$$

$$4 - 4 \cos^2 x + 9 \cos x - 6 = 0$$

$$4 \cos^2 x - 9 \cos x + 2 = 0$$

b $4 \cos^2 x - 9 \cos x + 2 = 0, 0 \leq \theta \leq 4\pi$

$$(4 \cos x - 1)(\cos x - 2) = 0$$

$\cos x = 2$ (no solution)

or $\cos x = 0.25 \Rightarrow x = 1.3, 5.0, 7.6, 11.2$

27 a $\tan 2x = 5 \sin 2x$

$$\frac{\sin 2x}{\cos 2x} = 5 \sin 2x$$

$$\sin 2x = 5 \sin 2x \cos 2x$$

$$(1 - 5 \cos 2x) \sin 2x = 0$$

b $\sin 2x(1 - 5 \cos 2x) = 0, 0 \leq x \leq \pi$

$$\text{Let } X = 2x$$


$$\sin X(1 - 5 \cos X) = 0, 0 \leq X \leq 2\pi$$

$$\sin X = 0 \Rightarrow X = 0, \pi, 2\pi$$

$$\text{or } \cos X = 0.2 \Rightarrow X = 1.37, 4.91$$

$$x = 0, 0.7, \frac{\pi}{2}, 2.5, \pi$$

28 a



b $\left(0, \frac{\sqrt{3}}{2}\right), \left(\frac{\pi}{3}, 0\right), \left(\frac{4\pi}{3}, 0\right)$

c $\cos\left(x + \frac{\pi}{6}\right) = 0.65, 0 \leq x \leq 2\pi$

$$\text{Let } X = x + \frac{\pi}{6}$$

$$\cos X = 0.65, \frac{\pi}{6} \leq X \leq \frac{13\pi}{6}$$

$$X = 0.863, 5.42$$

$$x = 0.34, 4.90$$

29 $\sin\left(3x + \frac{\pi}{3}\right) = 0.45, 0 \leq x \leq \pi$

Let $X = 3x + \frac{\pi}{3}, \frac{\pi}{3} \leq X \leq \frac{10\pi}{3}$

$\sin X = 0.45$

$X = 2.67, 6.75, 8.96$

$x = 0.54, 1.90 \text{ or } 2.64 \text{ (2 d.p.)}$

c $\sin 4\theta = 37 - 2 \cos 2\theta$

When θ is small:

$\text{LHS} \approx 4\theta$

and $\text{RHS} \approx 37 - 2\left(1 - \frac{(2\theta)^2}{2}\right)$

so $4\theta = 37 - 2 + 4\theta^2$

$4\theta^2 - 4\theta + 35 = 0$

$b^2 - 4ac < 0$

So there are no solutions.

Challenge

a $9 \sin \theta \tan \theta + 25 \tan \theta = 6$

When θ is small:

$\text{LHS} \approx 9\theta^2 + 25\theta$

so $9\theta^2 + 25\theta = 6$

$9\theta^2 + 25\theta - 6 = 0$

$(9\theta - 2)(\theta + 3) = 0$

$\theta = \frac{2}{9}$ or $\theta = -3$

$\theta = \frac{2}{9}$ is ‘small’, so this value is valid.

$\theta = -3$ is not ‘small’, so this value is not valid. ‘Small’ in this context is ‘close to 0’.

b $2 \tan \theta + 3 = 5 \cos 4\theta$

When θ is small:

$\text{LHS} \approx 2\theta + 3$ and $\text{RHS} \approx 5\left(1 - \frac{(4\theta)^2}{2}\right)$

so $2\theta + 3 = 5 - 40\theta^2$

$40\theta^2 + 2\theta - 2 = 0$

$20\theta^2 + \theta - 1 = 0$

$(4\theta + 1)(5\theta - 1) = 0$

$\theta = -\frac{1}{4}, \theta = \frac{1}{5}$

Both values of θ could be considered ‘small’ in this case so both solutions are valid.