

Trigonometric Functions 6D

1 a Use $1 + \tan^2 \theta = \sec^2 \theta$
with θ replaced with $\frac{1}{2}\theta$

$$1 + \tan^2\left(\frac{1}{2}\theta\right) = \sec^2\left(\frac{1}{2}\theta\right)$$

b $(\sec \theta - 1)(\sec \theta + 1)$ (multiply out)
 $= \sec^2 \theta - 1$
 $= (1 + \tan^2 \theta) - 1$
 $= \tan^2 \theta$

c $\tan^2 \theta (\cosec^2 \theta - 1)$
 $= \tan^2 \theta ((1 + \cot^2 \theta) - 1)$
 $= \tan^2 \theta \cot^2 \theta$
 $= \tan^2 \theta \times \frac{1}{\tan^2 \theta}$
 $= 1$

d $(\sec^2 \theta - 1)\cot \theta$
 $= \tan^2 \theta \cot \theta$
 $= \tan^2 \theta \times \frac{1}{\tan \theta}$
 $= \tan \theta$

e $(\cosec^2 \theta - \cot^2 \theta)^2$
 $= ((1 + \cot^2 \theta) - \cot^2 \theta)^2$
 $= 1^2 = 1$

f $2 - \tan^2 \theta + \sec^2 \theta$
 $= 2 - \tan^2 \theta + (1 + \tan^2 \theta)$
 $= 2 - \tan^2 \theta + 1 + \tan^2 \theta$
 $= 3$

g $\frac{\tan \theta \sec \theta}{1 + \tan^2 \theta}$
 $= \frac{\tan \theta \sec \theta}{\sec^2 \theta}$
 $= \frac{\tan \theta}{\sec \theta}$
 $= \tan \theta \cos \theta$
 $= \frac{\sin \theta}{\cos \theta} \times \cos \theta$
 $= \sin \theta$

h $(1 - \sin^2 \theta)(1 + \tan^2 \theta)$
 $= \cos^2 \theta \times \sec^2 \theta$
 $= \cos^2 \theta \times \frac{1}{\cos^2 \theta}$
 $= 1$

i $\frac{\cosec \theta \cot \theta}{1 + \cot^2 \theta}$
 $= \frac{\cosec \theta \cot \theta}{\cosec^2 \theta}$
 $= \frac{1}{\cosec \theta} \times \cot \theta$
 $= \frac{\sin \theta}{1} \times \frac{\cos \theta}{\sin \theta}$
 $= \cos \theta$

j $\sec^4 \theta - 2 \sec^2 \theta \tan^2 \theta + \tan^4 \theta$
 $= (\sec^2 \theta - \tan^2 \theta)^2$ (factorise)
 $= ((1 + \tan^2 \theta) - \tan^2 \theta)^2$
 $= 1^2 = 1$

k $4 \cosec^2 2\theta + 4 \cosec^2 2\theta \cot^2 2\theta$
 $= 4 \cosec^2 2\theta (1 + \cot^2 2\theta)$
 $= 4 \cosec^2 2\theta \cosec^2 2\theta$
 $= 4 \cosec^4 2\theta$

2 $\cosec x = \frac{k}{\cosec x}$
 $\Rightarrow \cosec^2 x = k$
 $\Rightarrow 1 + \cot^2 x = k$
 $\Rightarrow \cot^2 x = k - 1$
 $\Rightarrow \cot x = \pm \sqrt{k - 1}$

3 a $\cot \theta = \sqrt{3}$ $90^\circ < \theta < 180^\circ$

$$\Rightarrow \cot^2 \theta = 3$$

$$\Rightarrow 1 + \cot^2 \theta = 1 + 3 = 4$$

$$\Rightarrow \operatorname{cosec}^2 \theta = 4$$

$$\Rightarrow \sin^2 \theta = \frac{1}{4}$$

$$\Rightarrow \sin \theta = \frac{1}{2}$$

(as θ is in 2nd quadrant, $\sin \theta$ is positive)

b Using $\sin^2 \theta + \cos^2 \theta = 1$


$$\Rightarrow \cos^2 \theta = 1 - \sin^2 \theta = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\Rightarrow \cos \theta = -\frac{\sqrt{3}}{2}$$

(as θ is in 2nd quadrant, $\cos \theta$ is negative)

4 $\tan \theta = \frac{3}{4}$ $180^\circ < \theta < 270^\circ$

Draw a right-angled triangle where $\tan \varphi = \frac{3}{4}$



Using Pythagoras' theorem, $x = 5$

$$\text{So } \cos \varphi = \frac{4}{5} \text{ and } \sin \varphi = \frac{3}{5}$$

As θ is in the 3rd quadrant, both $\sin \theta$ and $\cos \theta$ are negative.

a $\sec \theta = \frac{1}{\cos \theta} = -\frac{1}{\cos \varphi} = -\frac{5}{4}$


b $\cos \theta = -\cos \varphi = -\frac{4}{5}$

c $\sin \theta = -\sin \varphi = -\frac{3}{5}$

5 $\cos \theta = \frac{24}{25}$, θ reflex

As $\cos \theta$ is positive and θ reflex, θ is in the 4th quadrant.

Use right-angled triangle where $\cos \varphi = \frac{24}{25}$



Using Pythagoras' theorem,

$$25^2 = x^2 + 24^2$$

$$\Rightarrow x^2 = 25^2 - 24^2 = 49$$

$$\Rightarrow x = 7$$

$$\text{So } \tan \varphi = \frac{7}{24} \text{ and } \sin \varphi = \frac{7}{25}$$

As θ is in the 4th quadrant, both $\tan \theta$ and $\sin \theta$ are negative

a $\tan \theta = -\frac{7}{24}$

b $\operatorname{cosec} \theta = \frac{1}{\sin \theta} = -\frac{1}{\frac{7}{25}} = -\frac{25}{7}$

6 a LHS $\equiv \sec^4 \theta - \tan^4 \theta$
 $\equiv (\sec^2 \theta - \tan^2 \theta)(\sec^2 \theta + \tan^2 \theta)$
(difference of two squares)
 $\equiv (1 + \tan^2 \theta)(\sec^2 \theta + \tan^2 \theta)$
(as $1 + \tan^2 \theta \equiv \sec^2 \theta$)
 $\Rightarrow \sec^2 \theta - \tan^2 \theta \equiv 1$
 $\equiv \sec^2 \theta + \tan^2 \theta \equiv \text{RHS}$

b LHS $\equiv \operatorname{cosec}^2 x - \sin^2 x$
 $\equiv (1 + \cot^2 x) - (1 - \cos^2 x)$
 $\equiv 1 + \cot^2 x - 1 + \cos^2 x$
 $\equiv \cot^2 x + \cos^2 x \equiv \text{RHS}$

6 c LHS $\equiv \sec^2 A(\cot^2 A - \cos^2 A)$

$$\begin{aligned} &\equiv \frac{1}{\cos^2 A} \left(\frac{\cos^2 A}{\sin^2 A} - \cos^2 A \right) \\ &\equiv \frac{1}{\sin^2 A} - 1 \equiv \operatorname{cosec}^2 A - 1 \\ &\quad (\text{use } 1 + \cot^2 \theta = \operatorname{cosec}^2 \theta) \\ &\equiv 1 + \cot^2 A - 1 \\ &\equiv \cot^2 A \equiv \text{RHS} \end{aligned}$$

d RHS $\equiv (\sec^2 \theta - 1)(1 - \sin^2 \theta)$

$$\begin{aligned} &\equiv \tan^2 \theta \times \cos^2 \theta \\ &\quad (\text{use } 1 + \tan^2 \theta \equiv \sec^2 \theta \text{ and} \\ &\quad \cos^2 \theta + \sin^2 \theta \equiv 1) \\ &\equiv \frac{\sin^2 \theta}{\cos^2 \theta} \times \cos^2 \theta \equiv \sin^2 \theta \\ &\equiv 1 - \cos^2 \theta \equiv \text{LHS} \end{aligned}$$

e LHS $\equiv \frac{1 - \tan^2 A}{1 + \tan^2 A} \equiv \frac{1 - \tan^2 A}{\sec^2 A}$

$$\begin{aligned} &\equiv \frac{1}{\sec^2 A} (1 - \tan^2 A) \\ &\equiv \cos^2 A \left(1 - \frac{\sin^2 A}{\cos^2 A} \right) \\ &\equiv \cos^2 A - \sin^2 A \\ &\equiv (1 - \sin^2 A) - \sin^2 A \\ &\equiv 1 - 2 \sin^2 A \equiv \text{RHS} \end{aligned}$$

f RHS $\equiv \sec^2 \theta \operatorname{cosec}^2 \theta$

$$\begin{aligned} &\equiv \sec^2 \theta (1 + \cot^2 \theta) \\ &\equiv \sec^2 \theta + \frac{1}{\cos^2 \theta} \times \frac{\cos^2 \theta}{\sin^2 \theta} \\ &\equiv \sec^2 \theta + \frac{1}{\sin^2 \theta} \\ &\equiv \sec^2 \theta + \operatorname{cosec}^2 \theta \equiv \text{LHS} \end{aligned}$$

Alternatively

$$\begin{aligned} \text{LHS} &\equiv \sec^2 \theta + \operatorname{cosec}^2 \theta \equiv \frac{1}{\cos^2 \theta} + \frac{1}{\sin^2 \theta} \\ &\equiv \frac{\sin^2 \theta + \cos^2 \theta}{\cos^2 \theta \sin^2 \theta} \equiv \frac{1}{\cos^2 \theta \sin^2 \theta} \\ &\equiv \sec^2 \theta \operatorname{cosec}^2 \theta \equiv \text{RHS} \end{aligned}$$

g LHS $\equiv \operatorname{cosec} A \sec^2 A$

$$\begin{aligned} &\equiv \operatorname{cosec} A (1 + \tan^2 A) \\ &\equiv \operatorname{cosec} A + \frac{1}{\sin A} \times \frac{\sin^2 A}{\cos^2 A} \\ &\equiv \operatorname{cosec} A + \frac{\sin A}{\cos^2 A} \\ &\equiv \operatorname{cosec} A + \frac{\sin A}{\cos A} \times \frac{1}{\cos A} \\ &\equiv \operatorname{cosec} A + \tan A \sec A \equiv \text{RHS} \end{aligned}$$

h LHS $\equiv (\sec \theta - \sin \theta)(\sec \theta + \sin \theta)$

$$\begin{aligned} &\equiv \sec^2 \theta - \sin^2 \theta \\ &\equiv (1 + \tan^2 \theta) - (1 - \cos^2 \theta) \\ &\equiv 1 + \tan^2 \theta - 1 + \cos^2 \theta \\ &\equiv \tan^2 \theta + \cos^2 \theta \equiv \text{RHS} \end{aligned}$$

7 $3 \tan^2 \theta + 4 \sec^2 \theta = 5$

$$\begin{aligned} &\Rightarrow 3 \tan^2 \theta + 4(1 + \tan^2 \theta) = 5 \\ &\Rightarrow 3 \tan^2 \theta + 4 + 4 \tan^2 \theta = 5 \\ &\Rightarrow 7 \tan^2 \theta = 1 \\ &\Rightarrow \tan^2 \theta = \frac{1}{7} \\ &\Rightarrow \cot^2 \theta = 7 \\ &\Rightarrow \operatorname{cosec}^2 \theta - 1 = 7 \\ &\Rightarrow \operatorname{cosec}^2 \theta = 8 \\ &\Rightarrow \sin^2 \theta = \frac{1}{8} \end{aligned}$$

As θ is obtuse (in the 2nd quadrant),
 $\sin \theta$ is positive.

$$\text{So } \sin \theta = \sqrt{\frac{1}{8}} = \frac{1}{2\sqrt{2}} = \frac{\sqrt{2}}{4}$$

8 a $\sec^2 \theta = 3 \tan \theta \quad 0^\circ \leq \theta \leq 360^\circ$

$$\Rightarrow 1 + \tan^2 \theta = 3 \tan \theta$$


$$\Rightarrow \tan^2 \theta - 3 \tan \theta + 1 = 0$$

$$\tan \theta = \frac{3 \pm \sqrt{5}}{2}$$

(equation does not factorise)

For $\tan \theta = \frac{3+\sqrt{5}}{2}$,


calculator value is 69.1° (3 s.f.)



Solutions are $69.1^\circ, 249^\circ$

For $\tan \theta = \frac{3-\sqrt{5}}{2}$,

calculator value is 20.9° (3 s.f.)



Solutions are $20.9^\circ, 201^\circ$

Set of solutions: $20.9^\circ, 69.1^\circ, 201^\circ, 249^\circ$ (3 s.f.)

b $\tan^2 \theta - 2 \sec \theta + 1 = 0 \quad -\pi \leq \theta \leq \pi$

$$\Rightarrow (\sec^2 \theta - 1) - 2 \sec \theta + 1 = 0$$


$$\Rightarrow \sec^2 \theta - 2 \sec \theta = 0$$

$$\Rightarrow \sec \theta (\sec \theta - 2) = 0$$

$\Rightarrow \sec \theta = 2$ (as $\sec \theta$ cannot be 0)

$$\Rightarrow \cos \theta = \frac{1}{2}$$

$$\Rightarrow \theta = -\frac{\pi}{3}, \frac{\pi}{3}$$



c $\operatorname{cosec}^2 \theta + 1 = 3 \cot \theta \quad -180^\circ \leq \theta \leq 180^\circ$


$$\Rightarrow (1 + \cot^2 \theta) + 1 = 3 \cot \theta$$

$$\Rightarrow \cot^2 \theta - 3 \cot \theta + 2 = 0$$

$$\Rightarrow (\cot \theta - 1)(\cot \theta - 2) = 0$$

$\Rightarrow \cot \theta = 1$ or $\cot \theta = 2$

$$\Rightarrow \tan \theta = 1 \text{ or } \tan \theta = \frac{1}{2}$$



$$\tan \theta = 1 \Rightarrow \theta = -135^\circ, 45^\circ$$

$$\tan \theta = \frac{1}{2} \Rightarrow \theta = -153^\circ, 26.6^\circ \text{ (3 s.f.)}$$

8 d $\cot \theta = 1 - \operatorname{cosec}^2 \theta \quad 0 \leq \theta \leq 2\pi$

$$\Rightarrow \cot \theta = 1 - (1 + \cot^2 \theta)$$

$$\Rightarrow \cot \theta = -\cot^2 \theta$$


$$\Rightarrow \cot^2 \theta + \cot \theta = 0$$

$$\Rightarrow \cot \theta (\cot \theta + 1) = 0$$

$$\Rightarrow \cot \theta = 0 \text{ or } \cot \theta = -1$$

For $\cot \theta = 0$ refer to graph: $\theta = \frac{\pi}{2}, \frac{3\pi}{2}$

For $\cot \theta = -1$, $\tan \theta = -1$



$$\text{So } \theta = \frac{3\pi}{4}, \frac{7\pi}{4}$$

$$\text{Set of solutions: } \frac{\pi}{2}, \frac{3\pi}{4}, \frac{3\pi}{2}, \frac{7\pi}{4}$$

e $3\sec \frac{1}{2}\theta = 2\tan^2 \frac{1}{2}\theta \quad 0 \leq \theta \leq 360^\circ$

$$\Rightarrow 3\sec \frac{1}{2}\theta = 2(\sec^2 \frac{1}{2}\theta - 1)$$

(use $1 + \tan^2 A \equiv \sec^2 A$ with $A = \frac{1}{2}\theta$)

$$\Rightarrow 2\sec^2 \frac{1}{2}\theta - 3\sec \frac{1}{2}\theta - 2 = 0$$

$$\Rightarrow (2\sec \frac{1}{2}\theta + 1)(\sec \frac{1}{2}\theta - 2) = 0$$

$$\Rightarrow \sec \frac{1}{2}\theta = -\frac{1}{2} \text{ or } \sec \frac{1}{2}\theta = 2$$

Only $\sec \frac{1}{2}\theta = 2$ applies as

$\sec A \leq -1$ or $\sec A \geq 1$

$$\Rightarrow \cos \frac{1}{2}\theta = \frac{1}{2}$$

As $0 \leq \theta \leq 360^\circ$ so $0 \leq \frac{1}{2}\theta \leq 180^\circ$

Calculator value is 60°

This is the only value in the interval.

$$\text{So } \frac{1}{2}\theta = 60^\circ$$

$$\Rightarrow \theta = 120^\circ$$

f $(\sec \theta - \cos \theta)^2 = \tan \theta - \sin^2 \theta \quad 0 \leq \theta \leq \pi$

$$\Rightarrow \sec^2 \theta - 2\sec \theta \cos \theta + \cos^2 \theta$$

$$= \tan \theta - \sin^2 \theta$$

$$\Rightarrow \sec^2 \theta - 2 + \cos^2 \theta = \tan \theta - \sin^2 \theta$$

$$\left(\sec \theta \cos \theta = \frac{1}{\cos \theta} \times \cos \theta = 1 \right)$$

$$\Rightarrow (1 + \tan^2 \theta) - 2 + (\cos^2 \theta + \sin^2 \theta) \\ = \tan \theta$$

$$\Rightarrow 1 + \tan^2 \theta - 2 + 1 = \tan \theta$$

$$\Rightarrow \tan^2 \theta - \tan \theta = 0$$

$$\Rightarrow \tan \theta (\tan \theta - 1) = 0$$

$$\Rightarrow \tan \theta = 0 \text{ or } \tan \theta = 1$$

$$\tan \theta = 0 \Rightarrow \theta = 0, \pi$$

$$\tan \theta = 1 \Rightarrow \theta = \frac{\pi}{4}$$

$$\text{Set of solutions: } 0, \frac{\pi}{4}, \pi$$

g $\tan^2 2\theta = \sec 2\theta - 1 \quad 0 \leq \theta \leq 180^\circ$

$$\Rightarrow \sec^2 2\theta - 1 = \sec 2\theta - 1$$

$$\Rightarrow \sec^2 2\theta - \sec 2\theta = 0$$

$$\Rightarrow \sec 2\theta (\sec 2\theta - 1) = 0$$

$$\Rightarrow \sec 2\theta = 0 \text{ (not possible)}$$

$$\text{or } \sec 2\theta = 1$$

$$\Rightarrow \cos 2\theta = 1 \quad 0 \leq 2\theta \leq 360^\circ$$

Refer to graph of $y = \cos \theta$


$$\Rightarrow 2\theta = 0^\circ, 360^\circ$$

$$\Rightarrow \theta = 0^\circ, 180^\circ$$

8 h $\sec^2 \theta - (1 + \sqrt{3}) \tan \theta + \sqrt{3} = 1$

for $0 \leq \theta \leq 2\pi$

$$\begin{aligned}\Rightarrow (1 + \tan^2 \theta) - (1 + \sqrt{3}) \tan \theta + \sqrt{3} &= 1 \\ \Rightarrow \tan^2 \theta - (1 + \sqrt{3}) \tan \theta + \sqrt{3} &= 0 \\ \Rightarrow (\tan \theta - \sqrt{3})(\tan \theta - 1) &= 0 \\ \Rightarrow \tan \theta &= \sqrt{3} \text{ or } \tan \theta = 1\end{aligned}$$



First answer for $\tan \theta = \sqrt{3}$ is $\frac{\pi}{3}$

Second solution is $\pi + \frac{\pi}{3} = \frac{4\pi}{3}$

First answer for $\tan \theta = 1$ is $\frac{\pi}{4}$

Second solution is $\pi + \frac{\pi}{4} = \frac{5\pi}{4}$

Set of solutions: $\frac{\pi}{4}, \frac{\pi}{3}, \frac{5\pi}{4}, \frac{4\pi}{3}$

9 a $\tan^2 k = 2 \sec k$

$$\begin{aligned}\Rightarrow (\sec^2 k - 1) &= 2 \sec k \\ \Rightarrow \sec^2 k - 2 \sec k - 1 &= 0 \\ \Rightarrow \sec k = \frac{2 \pm \sqrt{8}}{2} &= \frac{2 \pm 2\sqrt{2}}{2} = 1 \pm \sqrt{2}\end{aligned}$$

As $\sec k$ has no values between -1 and 1

$$\sec k = 1 + \sqrt{2}$$

b $\cos k = \frac{1}{1 + \sqrt{2}} = \frac{\sqrt{2} - 1}{(1 + \sqrt{2})(\sqrt{2} - 1)}$

$$= \frac{\sqrt{2} - 1}{2 - 1} = \sqrt{2} - 1$$

c Solutions of

$$\tan^2 k = 2 \sec k, \quad 0 \leq k \leq 360^\circ$$

are solutions of $\cos k = \sqrt{2} - 1$

Calculator solution is 65.5° (1 d.p.)

$$\Rightarrow k = 65.5^\circ, 360^\circ - 65.5^\circ$$

$$= 65.5^\circ, 294.5^\circ \text{ (1 d.p.)}$$

10 a As $a = 4 \sec x$

$$\Rightarrow \sec x = \frac{a}{4}$$

$$\Rightarrow \cos x = \frac{4}{a}$$

As $\cos x = b$

$$\Rightarrow b = \frac{4}{a}$$

b $c = \cot x$

$$\Rightarrow c^2 = \cot^2 x$$

$$\Rightarrow \frac{1}{c^2} = \tan^2 x$$

$$\Rightarrow \frac{1}{c^2} = \sec^2 x - 1$$

(use $1 + \tan^2 x \equiv \sec^2 x$)

$$\Rightarrow \frac{1}{c^2} = \frac{a^2}{16} - 1 \quad \left(\sec x = \frac{a}{4} \right)$$

$$\Rightarrow 16 = a^2 c^2 - 16c^2 \text{ (multiply by } 16c^2)$$

$$\Rightarrow c^2(a^2 - 16) = 16$$

$$\Rightarrow c^2 = \frac{16}{a^2 - 16}$$

11 a $x = \sec \theta + \tan \theta$

$$\frac{1}{x} = \frac{1}{\sec \theta + \tan \theta}$$

$$= \frac{\sec \theta - \tan \theta}{(\sec \theta + \tan \theta)(\sec \theta - \tan \theta)}$$

$$= \frac{\sec \theta - \tan \theta}{\sec^2 \theta - \tan^2 \theta}$$

$$= \sec \theta - \tan \theta$$

(as $1 + \tan^2 \theta \equiv \sec^2 \theta$)

$$\Rightarrow \sec^2 \theta - \tan^2 \theta \equiv 1$$

$$\mathbf{11 \ b} \quad x + \frac{1}{x} = \sec \theta + \tan \theta + \sec \theta - \tan \theta \\ = 2 \sec \theta$$

$$\Rightarrow \left(x + \frac{1}{x} \right)^2 = 4 \sec^2 \theta \\ \Rightarrow x^2 + 2x \times \frac{1}{x} + \frac{1}{x^2} = 4 \sec^2 \theta \\ \Rightarrow x^2 + \frac{1}{x^2} + 2 = 4 \sec^2 \theta$$

$$\mathbf{12} \quad 2 \sec^2 \theta - \tan^2 \theta = p$$

$$\Rightarrow 2(1 + \tan^2 \theta) - \tan^2 \theta = p \\ \Rightarrow 2 + 2 \tan^2 \theta - \tan^2 \theta = p \\ \Rightarrow \tan^2 \theta = p - 2 \\ \Rightarrow \cot^2 \theta = \frac{1}{p-2} \quad (p \neq 2)$$

$$\cosec^2 \theta = 1 + \cot^2 \theta = 1 + \frac{1}{p-2} \\ = \frac{(p-2)+1}{p-2} = \frac{p-1}{p-2}, \quad p \neq 2$$